

# Use of quality rapid diagnostic testing for safe blood transfusion in resource-limited settings

D. Mbanya<sup>1,2</sup>

1) Faculty of Medicine and Biomedical Sciences, University of Yaoundé I and 2) Yaoundé University Teaching Hospital, Yaounde, Cameroon

## Abstract

Blood safety in sub-Saharan Africa is jeopardized by multiple and diverse factors, including the predominance of high-risk family/replacement donors and the high prevalence of transfusion-transmissible infections (TTIs). Thus, stringent diagnostic strategies are vital. Western blotting is costly and technically demanding, and nucleic acid testing technologies, which have been reported to reliably reduce the rate of TTI, are not available in resource-limited settings. Therefore, there is a need for reliable and affordable testing alternatives in these settings. Rapid diagnostic testing has been widely adopted in developing countries, but, for effectiveness in blood safety, highly sensitive tests and the strict selection of low-risk blood donors are indispensable. Although the pre-serological window period remains a source of residual risk for transmission of TTIs during blood transfusion, the combination antigen–antibody rapid tests could contribute significantly to shortening the window period. Thus, despite its limitations, rapid diagnostic testing continues to contribute significantly to blood safety, as a cost-effective means of enhancing screening for TTIs and reducing their transmission in resource-limited rural settings.

**Keywords:** HBV, HCV, HIV, rapid diagnostic testing, resource-limited setting, syphilis, TTI

**Article published online:** 7 March 2013

*Clin Microbiol Infect* 2013; **19**: 416–421

**Corresponding author:** D. Mbanya, B.P. 8046, Yaoundé, Cameroon  
**E-mail:** [dmbanya1@yahoo.co.uk](mailto:dmbanya1@yahoo.co.uk)

## Introduction

Screening for blood-transmissible pathogens is a critical aspect of blood safety. Where effective screening programmes exist, the risk of transmission of transfusion-transmissible infections (TTIs) has been reported to progressively and significantly decrease [1,2]. This has been clearly demonstrated in western countries, where the most reliable screening methods are used, including nucleic acid amplification testing technologies, which allow for very early detection of various infectious agents [3]. Other confirmatory tests often used include western blot assays, line immunoassays (LIAs), recombinant immunoblot assays, indirect fluorescent antibody assays, and ELISAs. Unfortunately, these technologies are only feasible in major African cities, where external funding is obtainable.

Paradoxically, blood safety in these resource-limited settings (RLSs), and especially in sub-Saharan Africa (SSA), is compromised by several factors [4], including the highest prevalence of TTIs and the chronic shortage of reagents for their diagnosis. The major TTIs described include human

immunodeficiency virus (HIV), hepatitis B virus (HBV), hepatitis C virus (HCV), syphilis, and malaria. Hence, the WHO recommends that all blood for transfusion in RLSs be mandatorily screened for HIV, HBV, HCV, and syphilis [5], with either a combination of HIV antigen–antibody or HIV antibodies, hepatitis B surface antigen (HBsAg), a combination of HCV antigen–antibody or HCV antibodies, and specific treponemal antibodies, respectively [6]. Despite this, several countries are unable to screen all donated blood for one or more of these markers [5].

Another major issue concerns voluntary, non-remunerated and regular blood donors, who are reported to have the lowest rates of TTI [7]. However, in these RLSs, family/replacement and, in some cases, paid donors, with high risks for TTI, provide 70–90% of donations [4,8–10]. Although these family and replacement donors are more cost-effective for RLSs [8,11] and more easily manageable than a centralized ‘western model’ based on a 100% voluntary, non-remunerated and regular blood donor system [8,12], it must be emphasized that increased blood safety is only achievable with low-risk

blood donors. In 2010, Choudary reported a 1–2 per 1000 risk for blood recipients of contaminated blood of acquiring viral, bacterial or parasitic infections [13], even with very low pathogen load. Hence, it is vital to meticulously select safe blood donors and ensure the use of quality screening methods.

Even in RLSs, the use of affordable alternative screening methods such as rapid diagnostic testing will contribute significantly to blood safety, including in the more rural settings, where small numbers of daily tests (e.g. <20 samples/day) are performed. A recent study in Tete, in Mozambique, has clearly illustrated the need for and effectiveness of the consistent use of quality-assured rapid diagnostic tests (RDTs) in screening donated blood [14].

Rapid diagnostic testing is currently available for the screening of various pathogens, including HIV, HBV, HCV, syphilis and malaria infections, and newer technologies are being developed continuously. These are simple instrument-free assays that may be performed by personnel who have received very little training, and no laboratory facilities are required. The test kits are generally low in cost, with prices in the WHO bulk procurement scheme ranging from US \$0.40 to US\$2.00 per test. In addition, the kits are easy to store (room temperature, with no need for refrigeration), and require very short intervals (10–30 min) to provide results. These characteristics have rendered rapid diagnostic testing especially user-friendly in RLSs. Consequently, rapid diagnostic testing has been integrated into, and served as a point-of-care tool, in various healthcare programmes. Alternative testing strategies and algorithms have also been developed for pathogen screening and confirmation based on quality rapid diagnostic testing, both for transfusion and other services, especially where only few blood units are collected [15]. A typical scenario in SSA would be blood transfusion; this is often an urgent clinical decision requiring immediate results, which rapid diagnostic testing could provide within 30 min. Thus, rapid diagnostic testing is indispensable in such situations.

This review focuses on alerting clinicians and heads of transfusion services to the current pitfalls of transfusion services in RLSs, and the use of rapid diagnostic testing as an alternative for providing safe blood in these settings.

### Pitfalls in Blood Transfusion Services in RLSs

Blood transfusion services in RLSs, especially in SSA, are inadequate in all aspects of the blood safety chain, from policy development through blood donor selection to the appropriate use of blood products [16]. In 2006, only 22 of the 46 member states in the African Region of the WHO had developed policies for blood transfusion [17], and implementation has

been equally slow. The lack of national coordination of designated blood transfusion budgets, trained staff, donor recruitment programmes and national testing strategies, the lack of national guidelines for the clinical use of blood and the lack of haemovigilance and quality management programmes all contribute to hampering blood safety in these RLSs. In addition, the high disparity in funding systems and financing mechanisms is an issue of concern. Some RLSs have benefited from external funding, including the President's Emergency Plan for AIDS Relief (PEPFAR) and from the CDC, Atlanta, and hence may boast of reasonably reliable transfusion services, but the issue of sustainability post-funding remains.

### Performance and Limitations of Rapid diagnostic Testing in Blood Transfusion Safety in RLSs

#### Principles of rapid diagnostic testing

RDTs are based on different principles, of which four major types have been described (see Editorial). Many of these tests have an internal sample addition control that validates each test run. Whole blood, serum, plasma, and even finger-prick blood specimens, saliva or oral fluids have been used with these tests. Various test kits based on these principles are commercially available for use in RLSs, and lateral-flow immunochromatographic assays seem to be most current.

A few organizations, including the WHO and the US Food and Drug Administration, carry out evaluations of test kits based on criteria set by the WHO. They include: the speed with which results are obtained (<30 min); accuracy (sensitivity and specificity of >99% and >98%, respectively); minimal specimen volume; variability in specimen type (whole blood, plasma, and serum); ease of use (no specialized equipment); and stability of the reagents at ambient temperatures or between 2°C and 8°C [18]. Details of some currently available quality RDTs, based on these criteria, are shown in Table 1 [19–21]. For further reading on RDT, see Table 2.

#### RDTs for HIV

Many RDTs for HIV utilize the gp41 antigen as the target antigen for detecting HIV. Serological markers such as antibodies to HIV-1, including group O, to HIV-2, and to the p24 antigen, which appears early in the infection, are detected, as well as viral RNA when possible. Thus, the diagnostic tests contain antigens to specific HIV proteins of one group or subtype, hence may vary in performance, according to the [9] HIV strain being detected [22].

High rates of false-positive HIV results have been reported in several studies [23–25], usually because of the relatively low

**TABLE 1.** Some available quality rapid diagnostic test kits for transfusion-transmissible infections [19–21] (the selection of examples is based mainly on high sensitivity and/or specificity)

| Name                                   | Manufacturer                         | Pathogen | Sensitivity (%) | Specificity (%) | Performance evaluation criteria |
|--|--------------------------------------|----------|-----------------|-----------------|---------------------------------|
| Determine HIV1/2                       | Abbott Laboratories                  | HIV      | 100             | 99.6            | WHO [19]                        |
| HIV 1/2 STAT-PAK                       | ChemBio Diagnostic Systems           | HIV      | 99.7            | 99.3            | WHO [20]                        |
| HIV 1/2 STAT-PAK Dipstick              | ChemBio Diagnostic Systems           | HIV      | 99.0            | 100             | WHO [20]                        |
| Uni-Gold HIV-1/HIV-2                   | Trinity Biotech                      | HIV      | 100             | 100             | WHO [20]                        |
| Immunocomb II HIV 1&2 BiSpot           | Orgenics                             | HIV      | 100             | 99.7            | WHO [20]                        |
| Retrocheck HIV 1&2/Core HIV 1&2        | Qualpro Diagnostics/Core Diagnostics | HIV      | 100             | 99.1            | WHO [20]                        |
| DoubleCheckGold HIV 1&2                | Orgenics                             | HIV      | 100             | 99.3            | WHO [20]                        |
| OraQuick HIV-1/2                       | OraSure Technologies                 | HIV      | 100             | 99.2            | WHO [20]                        |
| Multispot HIV 1/2                      | Bio-Rad Laboratories                 | HIV      | 100             | 99.93           | FDA [19]                        |
| Determine Syphilis TP                  | Abbott Laboratories                  | Syphilis | 100             | 98.6            | WHO [19]                        |
| HCV Tri Dot                            | J. Mitra & Co.                       | HCV      | 100             | 91.5            | WHO [19]                        |
| HCV Spot                               | Genelabs Diagnostics                 | HCV      | 100             | 93.7            | WHO [19]                        |
| SeroCard HCV                           | Trinity Biotech                      | HCV      | 98.5            | 100             | WHO [19]                        |
| Determine HBsAg                        | Abbott Laboratories                  | HBV      | 100             | 100             | WHO [19]                        |
| Dainascreeen                           | Abbott Laboratories                  | HBV      | 100             | 100             | WHO [19]                        |
| SD BioLine HBsAg (One Step HBsAg Test) | Standard Diagnostics                 | HBsAg    | 97.95           | 100             | ICBS [21]                       |
| Assure HBsAg Rapid Test                | MP Biomedicals Asia Pacific          | HBsAg    | 97.95           | 100             | ICBS [21]                       |
| Quick Chaser HBsAg                     | Mizuho Medy                          | HBsAg    | 97.95           | 100             | ICBS [21]                       |

FDA, Food and Drug Administration; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; HCV, hepatitis C virus; HIV, human immunodeficiency virus; ICBS, International Consortium for Blood Safety.

**TABLE 2.** References on quality rapid diagnostic tests for further reading

| Pathogen                              | References   |
|---------------------------------------|--|
| HIV                                   | Point-of-Care HIV Testing Using Rapid HIV Test Kits: Guidance for Health-Care Professionals<br><a href="http://www.who.int/diagnostics_laboratory/evaluations/hiv/en/">www.who.int/diagnostics_laboratory/evaluations/hiv/en/</a><br>Rapid immunochromatographic detection by amplification of the colloidal gold signal<br><a href="http://www.who.int/diagnostics_laboratory/evaluations/hiv/en/">www.who.int/diagnostics_laboratory/evaluations/hiv/en/</a><br><a href="http://jcm.asm.org/content/48/8/2836.full.pdf+html">http://jcm.asm.org/content/48/8/2836.full.pdf+html</a>  |
| HIV<br><i>Trypanosoma</i><br>Syphilis | Sena AC, White BL, Sparling PF. Novel treponema pallidum serologic tests: a paradigm shift in syphilis screening for the 21st century. <i>Clin Infect Dis</i> 2010; 51: 700–708.<br><a href="http://cid.oxfordjournals.org/content/51/6/700.full">cid.oxfordjournals.org/content/51/6/700.full</a><br>Diagnostic testing for syphilis<br><a href="http://www.uptodate.com/contents/diagnostic-testing-for-syphilis">www.uptodate.com/contents/diagnostic-testing-for-syphilis</a><br>WHO Hepatitis C test kit evaluations<br><a href="http://www.who.int/diagnostics_laboratory/evaluations/hepc/en/">www.who.int/diagnostics_laboratory/evaluations/hepc/en/</a><br>Smith BD, Drobeniuc J, Jewett A et al. Evaluation of three rapid screening assays for detection of antibodies to hepatitis C virus. <i>J Infect Dis</i> 2011; 204: 825–831.<br><a href="http://www.natap.org/2011/HCV/JDis2011-Smith-825-31.pdf">www.natap.org/2011/HCV/JDis2011-Smith-825-31.pdf</a><br>O'Connell RJ, Gates RG, Bautista CT et al. Laboratory evaluation of rapid test kits to detect hepatitis C antibody for use in predonation screening in emergency settings. <i>Transfusion</i> 2012; Jul 23. doi: 10.1111/j.1537-2995.2012.03770.x. [Epub ahead of print]<br>CBS: HCV test kit (2006)<br><a href="http://www.icbs-web.org/page30.html">http://www.icbs-web.org/page30.html</a><br><a href="http://www.rapid-diagnostics.org/files/HBV-Manufacturers-website.rtf">http://www.rapid-diagnostics.org/files/HBV-Manufacturers-website.rtf</a><br><a href="http://www.who.int/diagnostics_laboratory/evaluations/hepb/en/">www.who.int/diagnostics_laboratory/evaluations/hepb/en/</a><br><a href="http://www.icbs-web.org/page30.html">http://www.icbs-web.org/page30.html</a> (2010) |
| HBsAg                                 |  |
| HIV, HCV, HBV, syphilis               | WHO 2010, <a href="http://www.who.int/bloodsafety/ScreeningDonatedBloodforTransfusion.pdf">http://www.who.int/bloodsafety/ScreeningDonatedBloodforTransfusion.pdf</a>  |

HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; HCV, hepatitis C virus; HIV, human immunodeficiency virus.

specificities of some test kits, despite very high sensitivities. Low specificities of HIV antibody screening tests may also result from tropical infections that cause B-cell activation, such as human African *Trypanosoma* infections, resulting in false-positive results [26]. False-negative results also represent a challenge [25], exacerbated by poor staff training and subjectivity in interpreting faint RDT results [24].

#### RDTs for HBV

The earliest HBV serological marker is its surface antigen HBsAg, but others develop in the course of the infection. Thus, one main target in HBV screening includes serology of HBsAg; however, routine screening for its core antibody (anti-HBc) is not recommended, because of non-specific results [6].

Most RDTs for HBV are based on agglutination or lateral-flow principles, and qualitatively detect the presence of HBsAg. These are also available for detecting other HBV serological markers, such as the HBe antigen (HBeAg) and antibodies to HBs, HBc and HBe. Various improvements are being made to these lateral-flow assays for HBV. As declared in one analysis in 2008 [27], 'the major challenge for HBsAg rapid tests is to detect the low levels of the target antigen that are present in a relatively high proportion of asymptomatic blood donors in order to achieve a clinical sensitivity similar to that of enzyme immune-assays (EIA)'. With the wide genetic diversity in HBV (genotypes A–H), there is high antigenic variation of HBsAg, and a need for high antigen concentrations of some variants in order for them to be detected with commercial kits. An

evaluation of various HBsAg assays using panels from the International Consortium on Blood Safety showed low analytical sensitivities of RDTs for HBsAg detection, as shown by 13 of 19 RDTs not detecting International Consortium on Blood Safety dilutions as positive, despite having high specificities [21]. Thus, sensitivity remains an issue for current HBV RDTs (HBsAg), despite excellent specificity. This emphasizes the continuous quest to improve the sensitivity of RDTs, which is indispensable for blood safety. An example of such an improvement is the new HBsAg rapid immunochromatographic assay based on the signal amplification system, which has been evaluated and shown to have enhanced sensitivity [27].

### RDTs for HCV

The laboratory diagnosis of HCV infection consists of detecting anti-HCV antibodies in the blood, followed by the confirmation of reactive samples with LIAs. Anti-HCV antibodies are detectable in approximately 90% of cases by 12 weeks of infection [28]. The number of available HCV antibody tests, including simple/rapid tests, has increased in recent years, but rapid diagnostic testing for HCV has been reported to be less sensitive than enzyme immunoassays (EIAs) [29,30]. A recent evaluation of anti-HCV RDT performance by Smith *et al.* [31] confirmed low sensitivities, but, interestingly, the recently US Food and Drug Administration-approved OraQuick HCV Rapid Antibody Test was reported to perform comparably to EIA techniques [32]. Nevertheless, serological screening has considerably reduced HCV transmission through blood transfusion [6].

### RDTs for screening for syphilis

Several methods are used for the diagnosis of syphilis; some require the use of specific treponemal antigens, some use rapid point-of-care tests, and others use EIAs as well as chemiluminescence assays. Non-treponemal tests are also used, including the old rapid plasma reagin test, which uses phospholipid (non-treponemal) antigens, the Venereal Disease Research Laboratory test, and the toluidine red unheated serum test, all of which are not only low in cost, but are also easy to perform. However, they require treponemal-based confirmation, because detectable antibodies can be produced by other inflammatory conditions, and their sensitivities vary with the type of test and stage of infection. Some treponemal tests include microhaemagglutination assays, particle agglutination assays, haemagglutination assays, and the fluorescent treponemal antibody absorption assay.

Several rapid syphilis tests are available worldwide that may be used with serum, plasma, or whole blood specimens [33]. These are highly sensitive and specific [33,34], as most detect both IgM and IgG. Although RDTs used in screening for syphilis

infection have been reported to be highly sensitive, they cannot distinguish between active and treated syphilis, so false-positive reactions are possible [35]. Thus, positive results need to be confirmed with quantitative non-treponemal testing to determine recent infection and response to therapy.

### Confirmation testing

The confirmation of RDT results is of relevance for issuing accurate results to blood donors, as well as for purposes of acquiring accurate epidemiological data; for blood safety, sensitivity is of the utmost importance. Where no established quality systems exist in RLSs, confirmation is not necessary, because all reactive blood units should always be discarded. However, in settings with an established quality system, the WHO recommends repeat testing, in duplicate, of the same sample and with the same assay before conclusions are drawn [6]. Furthermore, repeat testing may be performed with an alternative assay, either an RDT or an EIA [24,36]. Indeed, the WHO established an HIV diagnostic algorithm that has been adapted in various countries of RLSs for confirmation of antibodies to HIV. It must be noted here that confirmation rates are generally very low for HCV [37], although the rate of confirmation of most HBsAg RDT results is >90%, suggesting that confirmation with an alternative screening assay may not be necessary. More recently, a dual-path platform RDT based on the parallel detection of both treponemal and non-treponemal lines was evaluated in China, and showed reasonably high sensitivities and specificities with different specimen types (whole blood, plasma, and finger-prick blood), thus serving as both a screen and a confirmatory test [38]. This is a promising alternative for syphilis diagnosis in RLSs.

The main issue with TTI transmission is the window period, which allows for residual risks in blood transfusion, and combination antigen-antibody RDTs would shorten this [39], although they may still be costly for RLSs. In any case, very sensitive assays for TTIs are crucial for blood safety.

## Discussions and Conclusions

Today, western blotting remains costly and technically cumbersome for RLSs, and nucleic acid amplification testing remains inaccessible to and unaffordable for most countries, especially of SSA. In this light, testing strategies including simple rapid tests are indispensable, with the possibility of using serum, plasma, whole blood and even saliva in screening for different pathogens.

RDTs are not only cheaper than algorithms based on ELISA and western blot methods, but have also been reported to have comparable effectiveness [40,41]. Furthermore, a few

systematic evaluations of rapid HIV tests with non-B subtypes of HIV-1 group M and group O and HIV-2 have established that most tests adequately detect all subtypes of group M, but that their performance is more variable with group O and HIV-2 strains [22]. Interestingly, others have reported low levels of analytical sensitivity for RDTs [23,42]. In a field trial in Uganda, using an algorithm of three rapid HIV tests, of the weakly positive results on one or more of the three tests, 94.1% were negative or indeterminate on EIA or western blotting [23], further raising issues of sensitivity. Indeed, in one study in Cameroon, Tagny *et al.* demonstrated that using a combination antigen–antibody test could prevent 55 HIV transmissions per 10 000 donations of blood missed by RDTs [43]. In another study, conducted in KwaZulu Natal (South Africa), an increase was observed in the sensitivity of rapid diagnostic testing, from 68.7% to 93.5%, after a change of test brands [44]. Unfortunately, these variations in performance between RDTs are influenced by their quality, which depends on their costs; the lowest-performing assays tend to be low-cost, because they are neither evaluated nor approved by credible institutions. The lack of rigorous standards in the laboratory and untrained staff also contribute to inaccurate RDT results [44].

Recent WHO data indicated that, of 155 countries performing 100% screening for HIV, only 71 screened for HIV in a quality-assured manner [45]. Thus, quality assurance and quality control issues are other problems that continue to leave persistent gaps in the blood safety chain in RLSs. It is vital that quality assurance procedures be applied rigorously, and that procedures for detecting errors be included in all testing protocols, to maximize the accuracy of the laboratory results. Where possible, laboratories should participate in annual external quality assessments. Furthermore, the efforts to develop alternative, easily accessible and affordable, in-country-adapted technologies, including nucleic acid amplification testing technologies, should be fostered, to provide safe blood for all.

## Transparency Declaration

The author declares no conflicts of interest.

## References

- Pomper GJ, Wu Y, Snyder EL. Risks of transfusion-transmitted infections. *Curr Opin Hematol* 2003; 10: 412–418.
- Maresch C, Schluter PJ, Wilson AD, Sleight A. Residual infectious disease risk in screened blood transfusion from a high-prevalence population: Santa Catarina, Brazil. *Transfusion* 2008; 48: 273–281.
- Busch MP, Stramer SL, Kleinman SH. Evolving applications of nucleic acid amplification assays for prevention of virus transmission by blood components and derivatives. In: Garraty G, ed. *Applications of molecular biology in blood transfusion*. Bethesda, MD: American Association of Blood Banks, 1997, 123–176.
- Tayou Tagny C, Mbanya D, Tapko JB, Lefrere JJ. Blood safety in Sub-Saharan Africa: a multifactorial problem. *Transfusion* 2008; 48: 1256–1261.
- WHO. Fact Sheet No. 279. Blood safety and availability, WHO, June 2012. Available at: <http://www.who.int/mediacentre/factsheets/fs279/en/index.html>.
- World Health Organization. Screening donated blood for transfusion-transmissible infections. WHO, 2010. Available at: <http://www.who.int/bloodsafety/ScreeningDonatedBloodforTransfusion.pdf>.
- World Health Organization. WHO global database on blood safety, 2004–2005 report. Geneva, 2008.
- Medina LA, Kandulu J, Chisuwu L, Kashot A, Mundy C, Bates I. Laboratory costs of a hospital-based blood transfusion service in Malawi. *J Clin Pathol* 2007; 60: 1117–1120.
- World Health Organization. Blood safety and donation, Fact Sheet No. 279, WHO, June 2004. Available at: <http://www.who.int/mediacentre/factsheets/fs279/en/index.html>.
- Mbanya DN, Takam D, Ndumbe PM. Serological findings among first-time blood donors in Yaoundé, Cameroon: is safe donation a reality or a myth? *Transf Med* 2003; 13: 267–273.
- Bates I, Manyasi G, Medina LA. Reducing replacement donors in sub-Saharan Africa: challenges and affordability. *Transfus Med* 2007; 17: 434–442.
- Field SP, Allain JP. Transfusion in sub-Saharan Africa: does a Western model fit? *J Clin Pathol* 2007; 60: 1073–1075.
- Choudary N. Transfusion transmitted Infections: how many more? *Asian J Transfus Sci* 2010; 4: 71–72.
- Stokx J, Gillet P, De Weggheleire A *et al.* Seroprevalence of transfusion-transmissible infections and evaluation of the pre-donation screening performance at the Provincial Hospital of Tete, Mozambique. *BMC Infect Dis* 2011; 11: 141.
- Allain JP, Lee HH. Rapid tests for detection of viral markers in blood transfusion. *Expert Rev Mol Diagn* 2005; 5: 31–41.
- Tapko JB, Sam O, Diarra-Nama A. *Report on the status of blood safety in the WHO African Region for 2004*. Johannesburg: WHO Regional Office Africa, 2005.
- Tapko JB. The road to safe blood supply in the African region of the World Health Organization: trends and current status: 1999–2006. *Africa Sanguine* 2007; 10: 1–6.
- World Health Organization. *Laboratory biosafety manual*. Geneva: WHO, 1993.
- Davies J, van Oosterhout JJ, Nyirenda M *et al.* Reliability of rapid testing for hepatitis B in a region of high HIV endemicity. *Trans R Soc Trop Med Hyg* 2010; 104: 162–164.
- World Health Organization. List of HIV diagnostics eligible to tender for procurement by WHO in 2011. 2011. Available at: [http://www.who.int/diagnostics\\_laboratory/publications/evaluations/en/index.html](http://www.who.int/diagnostics_laboratory/publications/evaluations/en/index.html).
- Scheiblaue H, El-Nageh M, Diaz S *et al.* Performance evaluation of 70 hepatitis B virus (HBV) surface antigen (HBsAg) assays from around the world by a geographically diverse panel with an array of HBV genotypes and HBsAg subtypes. *Vox Sang* 2010; 98: 403–414.
- Phillips S, Granade TC, Pau CP, Candal D, Hu DJ, Parekh BS. Diagnosis of human immunodeficiency virus type 1 infection with different subtypes using rapid tests. *Clin Diagn Lab Immunol* 2000; 7: 698–699.
- Gray RH, Makumbi F, Serwadda D *et al.* Limitations of rapid HIV-1 tests during screening for trials in Uganda: diagnostic test accuracy study. *BMJ* 2007; 335: 188.
- Klarkowski DB, Wazome JM, Lokuge KM, Shanks L, Mills CF, O'Brien DP. The evaluation of a rapid in situ HIV confirmation test in a



- programme with a high failure rate of the WHO HIV two-test diagnostic algorithm. *PLoS ONE* 2009; 4: e4351.
25. Aghokeng AF, Mpoudi-Ngole E, Dimodi H *et al.* Inaccurate diagnosis of HIV-1 group M and O is a key challenge for ongoing universal access to antiretroviral treatment and HIV prevention in Cameroon. *PLoS ONE* 2009; 4: e7702.
  26. Lejon V, Mumba Ngoyi D, Ilunga M *et al.* Low specificities of HIV diagnostic tests caused by *Trypanosoma brucei gambiense* sleeping sickness. *J Clin Microbiol* 2010; 48: 2836–2839.
  27. Lin YH, Wang Y, Lou A *et al.* Evaluation of a new hepatitis B virus surface antigen rapid test with improved sensitivity. *J Clin Microbiol* 2008; 46: 3319–3324.
  28. Vallari DS, Jett BW, Alter HJ, Mimms LT, Holzman R, Shih JW. Serological markers of posttransfusion hepatitis C viral infection. *J Clin Microbiol* 1992; 30: 552–556.
  29. Scheiblaue H, El-Nageh M, Nick S, Fields H, Prince A, Diaz S. Evaluation of the performance of 44 assays used in countries with limited resources for the detection of antibodies to hepatitis C virus. *Transfusion*. 2006; 46: 708–718.
  30. Desbois D, Vaghefi P, Savary J, Dussaix E, Roque-Afonso AM. Sensitivity of a rapid immuno-chromatographic test for hepatitis C antibodies detection. *J Clin Virol* 2008; 41: 129–133.
  31. Smith BD, Teshale E, Jewett A *et al.* Performance of premarket rapid hepatitis C virus antibody assays in 4 national human immunodeficiency virus behavioral surveillance system sites. *Clin Infect Dis* 2011; 53: 780–786.
  32. Lee SR, Kardos KW, Schiff E *et al.* Evaluation of a new, rapid test for detecting HCV infection, suitable for use with blood or oral fluid. *J Virol Methods* 2011; 172: 27–31.
  33. World Health Organization. *The sexually transmitted diagnostics initiative (SDI): special programme for research and training in tropical diseases (TDR)*. Geneva: WHO, 2003.
  34. Herring AJ, Ballard RC, Pope V *et al.* A multi-center evaluation of nine, rapid, point-of-care syphilis tests using archived sera. *Sex Transm Infect* 2006; 82(suppl): v7–v12.
  35. van Dommelen L, Smismans A, Goossens VJ *et al.* Evaluation of a rapid one-step immunochromatographic test and two immunoenzymatic assays for the detection of anti-*Treponema pallidum* antibodies. *Sex Transm Infect* 2008; 84: 292–296.
  36. Constantine N. HIV InSite, 2006. Available at: [hivinsite.ucsf.edu/insite?page=kb-02-02-01](http://hivinsite.ucsf.edu/insite?page=kb-02-02-01) –
  37. Tulsiani S, Choudhury N, Desai P *et al.* True positivity of anti-hepatitis C virus enzyme-linked immunosorbent assay reactive blood donors: a prospective study done in western India. *Asian J Transfus Sci* 2012; 6: 165–168.
  38. Yin YP, Chen XS, Wei WH *et al.* A dual point-of-care test shows good performance in simultaneously detecting nontreponemal and treponemal antibodies in patients with syphilis: a multisite evaluation study in China. *Clin Infect Dis* 2013; 56: 659–665.
  39. Thoai Duong L, Laperche S, Brennan C. Evaluation of the sensitivity and specificity of six HIV combined p24 antigen and antibody assays. *J Virol Methods* 2004; 122: 185–194.
  40. Koblavi Demi S, Maurice C, Yavo D *et al.* Sensitivity and specificity of human immunodeficiency virus rapid serologic assays and testing algorithms in an antenatal clinic in Abidjan, Ivory Coast. *J Clin Microbiol* 2001; 39: 1808–1812.
  41. Urassa W, Nozohoor S, Jaffer S, Arama K, Mhalu F, Biberfeld G. Evaluation of an alternative confirmatory strategy for the diagnosis of HIV infection in Dar Es Salaam, Tanzania, based on simple rapid assays. *J Virol Methods* 2002; 100: 115–120.
  42. PATH. HealthTech historical profile: lateral-flow, point-of-care diagnostic tests for infectious disease, PATH, June 2005. Available at: <http://www.path.org/publications/details.php?i=1134>.
  43. Tagny CT, Mbanya D, Leballais L, Murphy E, Lefrere JJ, Laperche S. Reduction of the risk of transfusion-transmitted human immunodeficiency virus (HIV) infection by using an HIV antigen/antibody combination assay in blood donation screening in Cameroon. *Transfusion* 2011; 51: 184–190.
  44. Wolpaw BJ, Mathews C, Chopra M *et al.* The failure of routine rapid HIV testing: a case study of improving low sensitivity in the field. *BMC Health Serv Res* 2010; 10: 73.
  45. World Health Organization. *Blood safety indicators*, 2007. Geneva: WHO, 2009.